

ON THE FINITE GENERATION OF A FAMILY OF EXT MODULES

TONY J. PUTHENPURAKAL

Dedicated to Prof. L. L. Avramov on the occasion of his sixtieth birthday

ABSTRACT. Let Q be a Noetherian ring with finite Krull dimension and let $\mathbf{f} = f_1, \dots, f_c$ be a regular sequence in Q . Set $A = Q/(\mathbf{f})$. Let I be an ideal in A , and let M be a finitely generated A -module with $\text{projdim}_Q M$ finite. Set $\mathcal{R}(I) = \bigoplus_{n \geq 0} I^n$, the Rees-Algebra of I . Let $N = \bigoplus_{j \geq 0} N_j$ be a finitely generated graded $\mathcal{R}(I)$ -module. We show that

$$\bigoplus_{j \geq 0} \bigoplus_{i \geq 0} \text{Ext}_A^i(M, N_j)$$

is a finitely generated bi-graded module over $S = \mathcal{R}(I)[t_1, \dots, t_c]$. We give two applications of this result to local complete intersection rings.

1. INTRODUCTION

Let A be a Noetherian ring. Let I be an ideal in A and let M be a finitely generated A -module. M. Brodmann [4] proved that the set $\text{Ass}_A M/I^n M$ is independent of n for all large n . This result is usually deduced by proving that $\text{Ass}_A I^n M/I^{n+1} M$ is independent of n for all large n .

Some Generalizations of Brodmann's result

Fix $i \geq 0$. The following sets are independent of n for all large n .

- 1 (L. Melkerson and P. Schenzel) [10, Theorem 1]
 - (a) $\text{Ass}_A \text{Tor}_i^A(M, I^n/I^{n+1})$.
 - (b) $\text{Ass}_A \text{Tor}_i^A(M, A/I^n)$.
- 2 (same argument as in 1(a)).
 $\text{Ass}_A \text{Ext}_A^i(M, I^n/I^{n+1})$.
- 3 (D. Katz & E. West; [8, 3.5]) $\text{Ass}_A \text{Ext}_A^i(M, A/I^n A)$.

An example of A. Singh [12] shows that

$$\text{Ass}_A \lim_{\rightarrow} \text{Ext}_A^i(A/I^n, M) \quad \text{need not be finite.}$$

So in this example

$$\bigcup_{n \geq 1} \text{Ass}_A \text{Ext}_A^i(A/I^n, M) \quad \text{is not even finite.}$$

Date: September 12, 2008.

1991 Mathematics Subject Classification. Primary 13H10, 13D07; Secondary 13A02, 13A15.

Key words and phrases. local complete intersection, asymptotic associate primes, cohomological operators.

The work for this paper was done while the second author was visiting University of Kentucky by a fellowship from Department of Science and Technology, India. The author is deeply grateful to DST for its financial support and University of Kentucky for its hospitality.

We state some questions in this area which motivated me.

- (1) (W. Vasconcelos: [13, 3.5]) Is the set

$$\bigcup_{i \geq 0} \text{Ass}_A \text{Ext}_A^i(M, A) \quad \text{finite ?}$$

- (2) (L. Melkerson and P. Schenzel: [10, page 936]) Is the set

$$\bigcup_{i \geq 0} \bigcup_{n \geq 0} \text{Ass}_A \text{Tor}_i^A(M, A/I^n) \quad \text{finite ?}$$

The motivation for the main result of this paper came from a Vasconcelos's question. The author now does not believe that Vasconcelos's question has a positive answer in this generality. However he is unable to give a counter-example. Note that if A is a Gorenstein local ring then Vasconcelos's question has, trivially, a positive answer. If we change the question a little then we may ask: if M, D are two finitely generated A -modules then is the set

$$\bigcup_{i \geq 0} \text{Ass}_A \text{Ext}_A^i(M, D) \quad \text{finite ?}$$

This is not known for Gorenstein rings in general. Using Melkerson and Schenzel's question as a guidepost the questions I was interested to solve was

Let $(A, \mathfrak{m})$ be a local complete intersection of codimension c . Are the sets

- (a) $\bigcup_{i \geq 0} \bigcup_{j \geq 0} \text{Ass}_A \text{Ext}_A^i(M, D/I^j D) \quad \text{finite ?}$
 (b) $\bigcup_{i \geq 0} \bigcup_{j \geq 0} \text{Ass}_A \text{Ext}_A^i(M, I^j D) \quad \text{finite ?}$

In Theorem 5.9 we prove that (b) holds. I have been unable to verify whether (a) holds.

The main result in this paper is the following regarding finite generation of a family of Ext modules. Let $\mathcal{R}(I) = \bigoplus_{n \geq 0} I^n t^n$ be the Rees algebra of I .

Theorem 1. *Let Q be a Noetherian ring with finite Krull dimension and let $\mathbf{f} = f_1, \dots, f_c$ be a regular sequence in Q . Set $A = Q/(\mathbf{f})$. Let M be a finitely generated A -module with $\text{projdim}_Q M$ finite. Let I an ideal in A and let $N = \bigoplus_{n \geq 0} N_n$ be a finitely generated $\mathcal{R}(I)$ -module. Then*

$$\mathcal{E}(N) = \bigoplus_{i \geq 0} \bigoplus_{n \geq 0} \text{Ext}_A^i(M, N_n)$$

is a finitely generated bi-graded $\mathcal{S} = \mathcal{R}(I)[t_1, \dots, t_c]$ -module.

An easy consequence of this result is that (b) holds (by taking $N = \bigoplus_{n \geq 0} I^n D$); see Theorem 5.1. A complete local complete intersection ring is a quotient of a regular local ring mod a regular sequence. So in this case (b) holds from Theorem 5.1. The proof of (b) for local complete intersections in general is a little technical; see Theorem 5.9

We next discuss a surprising consequence of Theorem 1. Let $(A, \mathfrak{m})$ be a local complete intersection of codimension c . Let M, N be two finitely generated A -modules. Define

$$\text{cx}_A(M, N) = \inf \left\{ b \in \mathbb{N} \mid \varlimsup_{n \rightarrow \infty} \frac{\mu(\text{Ext}_A^n(M, N))}{n^{b-1}} < \infty \right\}$$

In this section 6 we prove, see Theorem 6.1, that

$$(\dagger) \quad \text{cx}_A(M, I^j N) \quad \text{is constant for all } j \gg 0.$$

We now describe in brief the contents of this paper. In section one we give a module structure to $\mathcal{E}(N)$ over $\mathcal{S}$ (as in Theorem 1). We also discuss a few preliminaries. The local case of Theorem 1 is proved in section 2 while the global case is proved in section 3. In section 4 we prove our results on asymptotic primes. In section 5 we prove $(\dagger)$.

Acknowledgements: The author thanks Prof. L. L. Avramov and Prof. J. Herzog for many discussions regarding this paper.

2. MODULE STRUCTURE

Let Q be a Noetherian ring and let $\mathbf{f} = f_1, \dots, f_c$ be a regular sequence in Q . Set $A = Q/(\mathbf{f})$. Let M be a finitely generated A -module with $\text{projdim}_Q M$ finite. *We will not change M throughout our discussion.* Let I an ideal in A and let $N = \bigoplus_{n \geq 0} N_n$ be a finitely generated $\mathcal{R}(I) = \bigoplus_{n \geq 0} I^n t^n$ -module. Set

$$\mathcal{E}(N) = \bigoplus_{i \geq 0} \bigoplus_{n \geq 0} \text{Ext}_A^i(M, N_n).$$

In this section we show $\mathcal{E}(N)$ is a bi-graded $\mathcal{S} = \mathcal{R}(I)[t_1, \dots, t_c]$ -module. We also discuss two preliminary results that we will need later in this paper.

2.1. Let $\mathbb{F} : \dots F_n \rightarrow \dots F_1 \rightarrow F_0 \rightarrow 0$ be a free resolution of M as a A -module.

Let $t_1, \dots, t_c : \mathbf{F}(+2) \rightarrow \mathbf{F}$ be the *Eisenbud-operators* [6, section 1.] Then

- (1) t_i are uniquely determined up to homotopy.
- (2) t_i, t_j commute up to homotopy.

2.2. Set $T = A[t_1, \dots, t_c]$ with $\deg t_i = 2$.

In [7] Gulliksen shows that if $\text{projdim}_Q M$ is finite then $\bigoplus_{i \geq 0} \text{Ext}_A^i(M, L)$ is a finitely generated T -module.

2.3. Let $N = \bigoplus_{n \geq 0} N_n$ be a f.g module over $\mathcal{R}(I)$. Let $u = xt^s$. The map

$$\begin{array}{ccc} N_n & \xrightarrow{u} & N_{n+1} \quad \text{yields} \\ \text{Hom}(\mathbf{F}, N_n) & \xrightarrow[t_r]{} & \text{Hom}(\mathbf{F}, N_n)(+2) \\ \downarrow u & & \downarrow u \\ \text{Hom}(\mathbf{F}, N_{n+s}) & \xrightarrow[t_r]{} & \text{Hom}(\mathbf{F}, N_{n+s})(+2) \end{array}$$

Taking homology gives that

$$\mathcal{E}(N) = \bigoplus_{i \geq 0} \bigoplus_{n \geq 0} \text{Ext}_A^i(M, N_n) \text{ is a bi-graded } \mathcal{S} = R(I)[t_1, \dots, t_c]\text{-module.}$$

Remark 2.4. 1. For each i , we have $\bigoplus_{n \geq 0} \text{Ext}_A^i(M, N_n)$ is a finitely generated $R(I)$ -module.

2. For each n , we have $\bigoplus_{i \geq 0} \text{Ext}_A^i(M, N_n)$ is a finitely generated $A[t_1, \dots, t_c]$ -module.

We state two Lemma's which will help us in proving Theorem 1.

2.5. Notation

(1) Let $N = \bigoplus_{n \geq 0} N_n$ be a graded $\mathcal{R}(I)$ -module. Fix $j \geq 0$. Set

$$N_{\geq j} = \bigoplus_{n \geq j} N_n.$$

$\mathcal{E}(N_{\geq j})$ is naturally isomorphic to the submodule

$$\mathcal{E}(N)_{\geq j} = \bigoplus_{i \geq 0} \bigoplus_{n \geq j} \mathcal{E}(N)_{ij}$$

of $\mathcal{E}(N)$.

(2) If $A \rightarrow A'$ is a ring extension and if D is an A -module then set $D' = D \otimes_A A'$. Notice that if D is finitely generated A -module then D' is a finitely generated A' -module.

(3) Set $\mathcal{S}' = \mathcal{S} \otimes_A A'$. Notice $\mathcal{S}'$ is a finitely generated bi-graded A' -algebra. Let $U = \bigoplus_{i \geq 0} \bigoplus_{n \geq 0} U_{i,n}$ be a graded $\mathcal{S}$ -module. Then

$$U' = U \otimes_A A' = \bigoplus_{i \geq 0} \bigoplus_{n \geq 0} U'_{i,n}$$

is a graded $\mathcal{S}'$ -module.

Lemma 2.6. *If $\mathcal{E}(N_{\geq j})$ is a finitely generated $\mathcal{S}$ -module then $\mathcal{E}(N)$ is a finitely generated $\mathcal{S}$ -module.*

Proof. Set $D = \mathcal{E}(N)/\mathcal{E}(N_{\geq j})$. We have the following exact sequence of $\mathcal{S}$ -modules

$$0 \longrightarrow \mathcal{E}(N_{\geq j}) \longrightarrow \mathcal{E}(N) \longrightarrow D \longrightarrow 0.$$

Using Gulliksen's result it follows that D is a finitely generated $T = A[t_1, \dots, t_c]$ -module. Since T is a subring of $\mathcal{S}$, we get that D is a finitely generated $\mathcal{S}$ -module. Thus if $\mathcal{E}(N_{\geq j})$ is a finitely generated $\mathcal{S}$ -module then $\mathcal{E}(N)$ is a finitely generated $\mathcal{S}$ -module $\square$

Lemma 2.7. *[with notation as in 2.5(3)] Let $A \rightarrow A'$ be a faithfully flat extension of rings and let $U = \bigoplus_{i \geq 0} \bigoplus_{n \geq 0} U_{i,n}$ be a graded $\mathcal{S}$ -module such that $U_{i,n}$ is a finitely generated A -module for each $i, n \geq 0$. If U' is a finitely generated $\mathcal{S}'$ -module. then U is a finitely generated $\mathcal{S}$ -module.*

Proof. Choose a finite generating set L of U' . Suppose $\deg_i u \leq a$ and $\deg_n u \leq b$ for each $u \in L$. For $0 \leq i \leq a$ and $0 \leq n \leq b$ choose a finite generating set $C_{i,n}$ of $U_{i,n}$. Set

$$C = \bigcup_{i=0}^a \bigcup_{n=0}^b C_{i,n}.$$

Set $C' = \{u \otimes 1 \mid u \in C\}$. Let V be the submodule of U generated by C . By construction C' generates U' . So $U' = V'$. Thus $(U/V) \otimes_A A' = 0$. Since A' is a faithfully flat A -algebra we get $U = V$. So U is a finitely generated $\mathcal{S}$ -module. $\square$

3. THE LOCAL CASE

In this section we prove Theorem 1 when $(Q, \mathfrak{n})$ is local. Let $\mathfrak{m}$ be the maximal ideal of A . Set $k = A/\mathfrak{m}$. Let I be an ideal in A . Set $F(I) = \mathcal{R}(I) \otimes_A k = \bigoplus_{n \geq 0} I^n/\mathfrak{m}I^n$ the fiber cone of I .

3.1. Assume $N = \bigoplus_{n \geq 0} N_n$ is a finitely generated $\mathcal{R}(I)$ -module. Notice

$$F(N) = N \otimes_A k = \bigoplus_{n \geq 0} N_n / \mathfrak{m} N_n$$

is a finitely generated $F(I)$ -module. Set

$$\text{spread}(N) := \dim_{F(I)} N / \mathfrak{m} N \quad \text{the analytic spread of } N.$$

Proof of Theorem 1 in the local case.

Case 1. *The residue field $k = A/\mathfrak{m}$ is infinite.*

We induct on $\text{spread}(N)$.

First assume $\text{spread}(N) = 0$. This implies that $N_n / \mathfrak{m} N_n = 0$ for all $n \gg 0$. By Nakayama Lemma, $N_n = 0$ for all $n \gg 0$; say $N_n = 0$ for all $n \geq j$. Then $\mathcal{E}(N_{\geq j}) = 0$ and it is obviously a finitely generated $\mathcal{S}$ -module. By 2.6 we get that $\mathcal{E}(N)$ is a finitely generated $\mathcal{S}$ -module.

When $\text{spread}(N) > 0$ then there exists $u = xt \in \mathcal{R}(I)_1$ which is $N \oplus F(N)$ -filter regular, i.e., there exists j such that

$$(0 :_N u)_n = 0 \quad \text{and} \quad (0 :_{F(N)} u)_n = 0 \quad \text{for all } n \geq j.$$

Set $N_{\geq j} = \bigoplus_{n \geq j} N_n$ and $U = N_{\geq j} / u N_{\geq j}$. Notice we have an exact sequence of $\mathcal{R}(I)$ -modules

$$0 \longrightarrow N_{\geq j}(-1) \xrightarrow{u} N_{\geq j} \longrightarrow U \longrightarrow 0.$$

For each $n \geq j$ the functor $\text{Hom}_A(M, -)$ induces the following long exact sequence of A -modules

$$\begin{aligned} 0 &\longrightarrow \text{Hom}_A(M, N_n) \xrightarrow{u} \text{Hom}_A(M, N_{n+1}) \longrightarrow \text{Hom}_A(M, U_{n+1}) \\ &\longrightarrow \text{Ext}_A^1(M, N_n) \xrightarrow{u} \text{Ext}_A^1(M, N_{n+1}) \longrightarrow \text{Ext}_A^1(M, U_{n+1}) \\ &\dots \quad \dots \quad \dots \quad \dots \quad \dots \quad \dots \quad \dots \quad \dots \quad \dots \\ &\longrightarrow \text{Ext}_A^i(M, N_n) \xrightarrow{u} \text{Ext}_A^i(M, N_{n+1}) \longrightarrow \text{Ext}_A^i(M, U_{n+1}) \\ &\dots \quad \dots \quad \dots \quad \dots \quad \dots \quad \dots \quad \dots \quad \dots \quad \dots \end{aligned}$$

Using the naturality of Eisenbud operators we have the following exact sequence of $\mathcal{S}$ -modules

$$\mathcal{E}(N_{\geq j})(-1, 0) \xrightarrow{(u, 0)} \mathcal{E}(N_{\geq j}) \longrightarrow \mathcal{E}(U)$$

By construction

$$\text{spread}(U) = \text{spread}(N_{\geq j}) - 1 = \text{spread}(N) - 1.$$

By induction hypothesis $\mathcal{E}(U)$ is a finitely generated $\mathcal{S}$ -module. Therefore by Lemma 3.2 we get $\mathcal{E}(N_{\geq j})$ is a finitely generated $\mathcal{S}$ -module. Using 2.6 we get that $\mathcal{E}(N)$ is finitely generated $\mathcal{S}$ -module.

Case 2. *The residue field k is finite.*

In this case we do the standard trick. Let $Q' = Q[X]_{\mathfrak{m}Q[X]}$. Set $A' = A \otimes_Q Q'$. Notice $A' = Q[X]_{\mathfrak{m}Q[X]}$ is a flat A -algebra with residue field $k(X)$ which is infinite. Set $I' = IA'$ and $M' = M \otimes_Q Q' = M \otimes_A A'$. Notice $\text{projdim}_{Q'} M'$ is finite. Set $\mathcal{R}(I)' = \mathcal{R}(I')$ the Rees algebra of I' . Notice that $N' = N \otimes_A A'$ is a finitely generated $\mathcal{R}(I)'$ -module. Also note that $\mathcal{E}(N') = \mathcal{E}(N) \otimes_A A'$.

By Case 1 we have that $\mathcal{E}(N')$ is a finitely generated $\mathcal{S}'$ -module. So by Lemma 2.7 we get that $\mathcal{E}(N)$ is a finitely generated $\mathcal{S}$ -module. $\square$

The next Lemma is a bi-graded version of Lemma 2.8 (1) from [11].

Lemma 3.2. *Let R be a Noetherian ring (not necessarily local) and let $B = \bigoplus_{i,j \geq 0} B_{i,j}$ be a finitely generated bi-graded R -algebra with $B_{0,0} = R$. Set*

$$B_x = \bigoplus_{i \geq 0} B_{(i,0)} \quad \text{and} \quad B_y = \bigoplus_{j \geq 0} B_{(0,j)}.$$

Let $V = \bigoplus_{i,j \geq 0} V_{i,j}$ be a bi-graded B -module such that

- (1) $V_{i,j}$ is a finitely generated R -module for each $i, j \geq 0$.
- (2) For each $i \geq 0$, $V_i = \bigoplus_{j \geq 0} V_{i,j}$ is finitely generated as a B_y -module
- (3) For each $j \geq 0$, $V_j = \bigoplus_{i \geq 0} V_{i,j}$ is finitely generated as a B_x -module
- (4) There exists $z \in B_{(r,0)}$ (with $r \geq 1$) such that we have the following exact sequence of B -modules

$$V(-r, 0) \xrightarrow{z} V \xrightarrow{\psi} D$$

where D is a finitely generated bi-graded B -module.

Then V is a finitely generated B -module

Proof. Step 1. We begin by reducing to the case when ψ is surjective.

Notice $D' = \text{image } \psi$ is a finitely generated bi-graded B -module. If $\psi' : V \rightarrow D'$ is the map induced by ψ then we have an exact sequence

$$V(-r, 0) \xrightarrow{z} V \xrightarrow{\psi'} D' \longrightarrow 0.$$

Thus we may assume ψ is surjective.

Step 2. Choosing generators:

2.1: Choose a finite set W in V of homogeneous elements such that

$$\psi(W) = \{\psi(w) \mid w \in W\}$$

is a generating set for D .

2.2: Assume all the elements in W have x -co-ordinate $\leq c$.

2.3: For each $i \geq 0$, by hypothesis V_i is a finitely generated B_y -module. So we may choose a finite set P_i of homogeneous elements in V_i which generates V_i as a B_y -module.

2.4: Set

$$G = W \cup \left(\bigcup_{i=0}^c P_i \right).$$

Clearly G is a finite set.

Claim: G is a generating set for V .

Let U be the B -submodule of V generated by G . It suffices to prove that $U_{i,j} = V_{i,j}$ for all $i, j \geq 0$. By construction we have that for $0 \leq i \leq c$

$$(*) \quad U_{i,j} = V_{i,j} \quad \text{for each } j \geq 0$$

Let $\preceq$ be the *lex-order* on $X = \mathbb{Z}_{\geq 0} \times \mathbb{Z}_{\geq 0}$. It is well-known that $\preceq$ is a total order on X . So we can prove our result by induction on X with respect to the total order $\preceq$.

The base case is $(0, 0)$.

In this case $U_{0,0} = V_{0,0}$ by $(*)$.

Let $(i, j) \in X \setminus \{(0, 0)\}$ and assume that for all $(r, s) \prec (i, j)$; we have $U_{i,j} = V_{i,j}$.

Subcase 1. $i \leq c$.

By $(*)$ we have $U_{i,j} = V_{i,j}$.

Subcase 2. $i > c$.

Let $p \in V_{i,j}$. By construction, note that there exists $w_1, \dots, w_m \in W \subseteq C$ such that

$$\psi(p) = \sum_{l=0}^m h_l \psi(w_l) \quad \text{where } h_i \in B.$$

We may assume that $\deg h_l w_l = (i, j)$ for each l . Set $p' = \sum_{i=0}^m h_i w_i \in V_{i,j}$. Notice

1. $p' \in U_{i,j}$.
2. $p - p' \in \ker \psi$.

So

$$p - p' = z \bullet q \quad \text{where } q \in V_{(i-r,j)}.$$

If $q = 0$ then $p = p' \in U_{i,j}$.

Otherwise note that $(i-r, j) \prec (i, j)$. So by induction hypothesis $q \in U_{(i-r,j)}$. It follows that $p \in U_{i,j}$.

Thus $V_{i,j} \subseteq U_{i,j}$. Since $U_{i,j} \subseteq V_{i,j}$ by construction it follows that $U_{i,j} = V_{i,j}$. The result follows by induction on X . $\square$

4. THE GLOBAL CASE:

We need quite a few preliminaries to prove the global case. See 4.2 for the difficulty in going from local to the global case. Note that in the local case we proved the result by inducting on $\text{spread}(N)$. This is unavailable to us in the global situation as there are usually infinitely many maximal ideals in a global ring. Most of this section we will discuss two invariants of a graded $\mathcal{R}(I)$ -module $N = \bigoplus_{n \geq 0} N_n$. We will use these invariants to prove Theorem 1 by induction.

4.1. Notation and Conventions: We take dimension of the zero module to be -1 . We define the zero-polynomial to have degree -1 .

Let $\mathfrak{P} \in \text{Spec } Q$. If $\mathfrak{P} \supseteq \mathfrak{f}$ then set $\mathfrak{p} = \mathfrak{P}/\mathfrak{f}$. If $\mathfrak{P} \not\supseteq \mathfrak{f}$ then any A -module localized at $\mathfrak{P}$ is zero. So assume $\mathfrak{P} \supseteq \mathfrak{f}$. Notice

- (1) $\mathcal{R}(I)_{\mathfrak{p}} \cong \mathcal{R}(IA_{\mathfrak{p}})$ and $\mathcal{S}_{\mathfrak{p}} \cong \mathcal{R}(I)_{\mathfrak{p}}[t_1, \dots, t_c]$
- (2) $M_{\mathfrak{p}} = M_{\mathfrak{P}}$ has finite projective dimension as a $Q_{\mathfrak{P}}$ -module.
- (3) $\mathcal{E}(N)_{\mathfrak{p}} \cong \mathcal{E}(N_{\mathfrak{p}})$.

4.2. The difficulty in going from local to global:

For each $\mathfrak{p} \in \text{Spec } A$ it follows from 4.1 that $\mathcal{E}(N)_{\mathfrak{p}}$ is a finitely generated $A_{\mathfrak{p}}$ -module. Usually $\text{Supp}_A \mathcal{E}(N)$ will be an infinite set. So we cannot apply the local case and conclude.

The situation when $\text{Supp}_A \mathcal{E}(N)$ is a finite set will help in the base step of our induction argument to prove Theorem 1. So we show it separately.

Lemma 4.3. *If $\text{Supp}_A \mathcal{E}(N)$ is a finite set then $\mathcal{E}(N)$ is a finitely generated $\mathcal{S}$ -module.*

Proof. We may choose a finite subset C of $\mathcal{E}(N)$ such that its image in $\mathcal{E}(N)_{\mathfrak{p}}$ generates $\mathcal{E}(N)_{\mathfrak{p}}$ for each $\mathfrak{p} \in \text{Supp}_A \mathcal{E}(N)$. Set U to be the finitely generated submodule of $\mathcal{E}(N)$ generated by C .

Set $D = \mathcal{E}(N)/U$. Notice that $D_{\mathfrak{p}} = 0$ for each $\mathfrak{p} \in \text{Spec } A$. So $D = 0$. Therefore $\mathcal{E}(N) = U$ is a finitely generated A -module. $\square$

4.4. First Inductive device:

Since N is a finitely generated $\mathcal{R}(I)$ -module we have $\text{ann}_A N_i \subseteq \text{ann}_A N_{i+1}$ for all $i \gg 0$. Since A is Noetherian it follows that $\text{ann}_A N_n$ is constant for all $n \gg 0$. Call this stable value $\mathfrak{L}_N$. This enables us to define *Limit dimension* of N .

$$\lim\text{Dim } N = \lim_{n \rightarrow \infty} \dim_A N_n = \dim A/\mathfrak{L}_N.$$

Since A has finite Krull-dimension we get that $\lim\text{Dim } N$ is finite.

4.5. Let $\mathfrak{P}$ be a prime ideal in A . If D is a finitely generated A -module then

$$\text{ann}_{A_{\mathfrak{P}}} D_{\mathfrak{P}} = (\text{ann}_A D)_{\mathfrak{P}} = (\text{ann}_A D) A_{\mathfrak{P}}.$$

Therefore

$$(\mathfrak{L}_N)_{\mathfrak{P}} = \mathfrak{L}_{N_{\mathfrak{P}}}.$$

4.6. Note that if $\lim\text{Dim}(N) = -1$ then $N_j = 0$, say for all $j \geq j_0$. So $\mathcal{E}(N_{\geq j_0}) = 0$. Using 2.6 it follows that $\mathcal{E}(N)$ is a finitely generated $\mathcal{S}$ -module. The first non-trivial case is the following

Proposition 4.7. *If $\lim\text{Dim}(N) = 0$ then $\mathcal{E}(N)$ is a finitely generated A -module.*

Proof. This implies that $A/\mathfrak{L}_N$ is Artinian. Say $\dim N_n = 0$ for $n \geq r$. Clearly

$$\text{Supp}_A \mathcal{E}(N_{\geq r}) \subseteq \text{Supp}_A A/\mathfrak{L}_N \quad \text{a finite set of maximal ideals in } A.$$

It follows from 4.3 that $\mathcal{E}(N_{\geq r})$ is a finitely generated $\mathcal{S}$ -module. Using 2.6 we get that $\mathcal{E}(N)$ is a finitely generated $\mathcal{S}$ -module. $\square$

4.8. higher degree filter-regular element

We do not have filter regular elements of degree 1 in the global situation. However we can do the following:

Set $E = N/H_{R_+}^0(N)$. Assume $E \neq 0$. As $H_{R_+}^0(E) = 0$ there exists homogeneous $u \in R_+$ such that u is E -regular, [5, 1.5.11]. Say $\deg u = s$. Since $E_n = N_n$ for all $n \gg 0$ it follows that the map $N_i \rightarrow N_{i+s}$ induced by multiplication by u is injective for all $i \gg 0$. We will say that u is a N filter-regular element of degree s .

4.9. The second inductive device:

We now discuss a global invariant of N which patches well with local ones.

4.10. The local invariant

Let $(A, \mathfrak{m})$ be local and let $W = \bigoplus_{n \geq 0} W_n$ be a finitely generated $\mathcal{R}(I)$ -module. For convenience we assume that $\mathfrak{L}_W = \text{ann}_A W_n$ for all $n \geq 0$. Let $\mathfrak{a} \subseteq \mathfrak{L}_W$ be an ideal. Fix $j \geq 0$. Set

$$d_{\mathfrak{a}}(W, j) = \begin{cases} 0, & \text{if } \dim W_j < \dim A/\mathfrak{a}, \\ e(\mathfrak{m}, W_j); & \text{otherwise.} \end{cases}$$

Note that W_j is an $A/\mathfrak{a}$ -module and that $d_{\mathfrak{a}}(W, j)$ is the modified multiplicity function on the $A/\mathfrak{a}$ -module W_j .

Remark 4.11. Notice if $\dim W_j = \dim A/\mathfrak{a}$ then

$$d_{\mathfrak{a}}(W, j) = d_{\mathfrak{L}_W}(W, j).$$

Let $\mu(D)$ denote the minimal number of generators of an A -module D .

Lemma 4.12. *The function $d_{\mathfrak{a}}(W, -)$ is polynomial of degree $\leq \mu(I) - 1$.*

Proof. We may assume that the residue field of A is infinite. Set $T = \mathcal{R}(I)/\mathfrak{a}\mathcal{R}(I) = \bigoplus_{n \geq 0} T_n$. Notice $T_0 = A/\mathfrak{a}$. Let $\mathbf{x} = x_1, \dots, x_r$ be a minimal reduction of $\mathfrak{m}(A/\mathfrak{a})$. So $e(\mathfrak{m}, -) = e(\mathbf{x}, -)$, cf. [5, 4.6.5]. Then by a result due to Serre, cf., [5, 4.7.6], we get that

$$e(\mathbf{x}, W_j) = \sum_{i=0}^r (-1)^i \ell(H_i(\mathbf{x}, W_j))$$

Notice $H_i(\mathbf{x}, W) = \bigoplus_{j \geq 0} H_i(\mathbf{x}, W_j)$ is a finitely generated $T/\mathbf{x}T$ -module. Notice $(T/\mathbf{x}T)_0 = A/(\mathfrak{a} + \mathbf{x})$ is Artinian. Furthermore $(T/\mathbf{x}T)_1$ is a quotient of $\mathcal{R}(I)_1$ and so can be generated by $\mu(I)$ elements. Therefore the function $j \mapsto \ell(H_i(\mathbf{x}, W_j))$ is polynomial of degree $\leq \mu(I) - 1$. The result follows. $\square$

Definition 4.13. $\theta(\mathfrak{a}, W)$ = degree of the polynomial function $d_{\mathfrak{a}}(W, -)$.

Clearly $\theta(\mathfrak{a}, W)$ is non-negative if and only if $\lim \dim W = \dim R/\mathfrak{a}$ and is -1 otherwise.

4.14. The global invariant

Let A be a Noetherian ring with finite Krull dimension. Let $I = (x_1, \dots, x_s)$ be an ideal in A . Let $W = \bigoplus_{n \geq 0} W_n$ be a finitely generated $\mathcal{R}(I)$ -module. For convenience we assume that $\mathfrak{L}_W = \text{ann}_A W_n$ for all $n \geq 0$. Let $\mathfrak{a} \subseteq \mathfrak{L}_W$ be an ideal. Set

$$\mathcal{C}(\mathfrak{a}) = \{\mathfrak{m} \mid \mathfrak{m} \in \text{m-Spec}(A), \mathfrak{m} \supseteq \mathfrak{a} \text{ \& \; } \dim(A/\mathfrak{a})_{\mathfrak{m}} = \dim A/\mathfrak{a}\}.$$

Let $I = (x_1, \dots, x_s)$. If $\mathfrak{m} \in \mathcal{C}(\mathfrak{a})$ then note that

- (a) $W_{\mathfrak{m}} = \bigoplus_{n \geq 0} (W_n)_{\mathfrak{m}}$.
- (b) $\mathfrak{L}_{W_{\mathfrak{m}}} = (\mathfrak{L}_W)_{\mathfrak{m}}$. So $\mathfrak{a}_{\mathfrak{m}} \subseteq \mathfrak{L}_{W_{\mathfrak{m}}}$
- (c) $\theta(\mathfrak{a}_{\mathfrak{m}}, W_{\mathfrak{m}}) \leq s - 1$.

Define

$$\theta(\mathfrak{a}, W) = \max\{\theta(\mathfrak{a}_{\mathfrak{m}}, W_{\mathfrak{m}}) \mid \mathfrak{m} \in \mathcal{C}(\mathfrak{a})\}.$$

By (c) above we get that $\theta(\mathfrak{a}, W)$ is defined and is $\leq s - 1$.

4.15. Properties of $\theta(\mathfrak{a}, W)$.

We describe some properties of $\theta(\mathfrak{a}, W)$ we need for the proof of global case of Theorem 1. Let $I = (x_1, \dots, x_s)$.

- (i) $\theta(\mathfrak{a}, W) \leq s - 1$.

This is clear.

- (ii) If $\mathfrak{L}_W \neq A$ then $\theta(\mathfrak{L}_W, W) \geq 0$.

It suffices to consider the local case. Note that then $d_{\mathfrak{L}_W}(W, j) > 0$ for all $j \geq 0$. It follows that $\theta(\mathfrak{L}_W, W) \geq 0$.

- (iii) $\theta(\mathfrak{a}, W) = -1$ if and only if $\lim \dim W < \dim A/\mathfrak{a}$.

This follows from the following four facts:

- (a) By definition of $\mathcal{C}(\mathfrak{a})$ we have that

$$\dim A/\mathfrak{a} = \dim(A/\mathfrak{a})_{\mathfrak{m}} \text{ for each } \mathfrak{m} \in \mathcal{C}(\mathfrak{a}).$$

- (b) If $\theta(\mathfrak{a}, W) = -1$ then $\theta(\mathfrak{a}_{\mathfrak{m}}, W_{\mathfrak{m}}) = -1$ for all $\mathfrak{m} \in \mathcal{C}(\mathfrak{a})$. This is equivalent to saying that $\lim \dim W_{\mathfrak{m}} < \dim(A/\mathfrak{a})_{\mathfrak{m}}$ for all $\mathfrak{m} \in \mathcal{C}(\mathfrak{a})$

- (c) Note that since $\mathfrak{a} \subseteq \mathfrak{L}_W$ we get that

$$\lim \dim W = \max\{\lim \dim W_{\mathfrak{m}} \mid \mathfrak{m} \in \mathcal{C}(\mathfrak{a})\}$$

(iv) If $\theta(\mathfrak{a}, W) \geq 0$ then $\theta(\mathfrak{L}_W, W) \leq \theta(\mathfrak{a}, W)$.

By previous item we get that $\limDim W = \dim A/\mathfrak{a}$. By hypothesis we also have $\mathfrak{a} \subseteq \mathfrak{L}_W$. Since $\dim A/\mathfrak{a} = \dim A/\mathfrak{L}_W$ it follows that $\mathcal{C}(\mathfrak{L}_W) \subseteq \mathcal{C}(\mathfrak{a})$. Using 4.11 it follows that $\theta(\mathfrak{L}_W, W) \leq \theta(\mathfrak{a}, W)$.

(v) $u \in \mathcal{R}(I)_+$ be homogeneous of degree b . Assume u is W -filter regular. Set $E = W/uW$. Then

$$\theta(\mathfrak{L}_W, E) \leq \theta(\mathfrak{L}_W, W) - 1.$$

Suppose $\theta(\mathfrak{L}_W, E) = \theta((\mathfrak{L}_W)_{\mathfrak{p}}, E_{\mathfrak{p}})$ for some $\mathfrak{p} \in \mathcal{C}(\mathfrak{a})$. Since u is W filter regular; multiplication by u induces the following exact sequence

$$0 \rightarrow W_{j-b} \rightarrow W_j \rightarrow E_j \rightarrow 0 \quad \text{for all } j \gg 0.$$

Localization at $\mathfrak{p}$ yields an exact sequence

$$0 \rightarrow (W_{j-b})_{\mathfrak{p}} \rightarrow (W_j)_{\mathfrak{p}} \rightarrow (E_j)_{\mathfrak{p}} \rightarrow 0 \quad \text{for all } j \gg 0.$$

Since $d_{\mathfrak{L}_W, \mathfrak{p}}(-, -)$ is an additive functor on $(A/\mathfrak{L}_W)_{\mathfrak{p}}$ -modules we get that

$$\theta((\mathfrak{L}_W)_{\mathfrak{p}}, E_{\mathfrak{p}}) = \theta((\mathfrak{L}_W)_{\mathfrak{p}}, W_{\mathfrak{p}}) - 1.$$

The result follows since

$$\begin{aligned} \theta((\mathfrak{L}_W)_{\mathfrak{p}}, E_{\mathfrak{p}}) &= \theta(\mathfrak{L}_W, E) \\ \text{and } \theta((\mathfrak{L}_W)_{\mathfrak{p}}, W_{\mathfrak{p}}) &\leq \theta(\mathfrak{L}_W, W). \end{aligned}$$

We now give a proof of our main result

Proof of Theorem 1. We induct on $\limDim N$.

If $\limDim N = -1, 0$ then the result follows from 4.7.

Assume $\limDim N \geq 1$ and assume the result holds for all $\mathcal{R}(I)$ -modules E with $\limDim E \leq \limDim N - 1$. Let $x \in \mathcal{R}(I)_+$ be homogeneous and a N -filter regular element. Set $D = N/xD$. By Lemma 2.6 it suffices to assume the case when x is N -regular.

We now induct on $\theta(\mathfrak{L}_N, N)$ If $\theta(\mathfrak{L}_N, N) = 0$ then $\theta(\mathfrak{L}_N, D) \leq -1$, by 4.15(v). Using 4.15(iii) we get that

$$\limDim D < \dim A/\mathfrak{L}_N = \limDim N.$$

By induction hypothesis (on $\limDim$) the module $\mathcal{E}(D)$ is finitely generated $\mathcal{S}$ -module. The short exact sequence of $\mathcal{R}(I)$ -modules

$$0 \rightarrow N(-r) \xrightarrow{x} N \rightarrow D \rightarrow 0$$

induces an exact sequence of $\mathcal{S}$ -modules

$$\mathcal{E}(N)(-r) \xrightarrow{x} \mathcal{E}(N) \rightarrow \mathcal{E}(D).$$

By Lemma 3.2 we get that $\mathcal{E}(N)$ is a finitely generated $\mathcal{S}$ -module.

We assume the result if $\theta(\mathfrak{L}_N, N) \leq i$ and prove when $\theta(\mathfrak{L}_N, N) = i + 1$. Let D be as above. So $\theta(\mathfrak{L}_N, D) \leq i$, by 4.15(v).

If $\theta(\mathfrak{L}_N, D) = -1$ then the argument as above yields $\mathcal{E}(N)$ to be a finitely generated $\mathcal{S}$ -module.

If $\theta(\mathfrak{L}_N, D) \geq 0$ then by 4.15(iv) we get that $\theta(\mathfrak{L}_D, D) \leq \theta(\mathfrak{L}_N, D) \leq i$. So by induction hypothesis on $\theta(-, -)$ we get that $\mathcal{E}(D)$ is a finitely generated $\mathcal{S}$ -module. By an argument similar to the above we get that $\mathcal{E}(N)$ is a finitely generated $\mathcal{S}$ -module. $\square$

5. APPLICATION I

ASYMPTOTIC ASSOCIATED PRIMES

In this section we give a proof of our main motivating question Theorem 5.9. We also give two proofs of Theorem 5.1.

Theorem 5.1. *Let Q be a Noetherian ring with finite Krull dimension and let $\mathbf{f} = f_1, \dots, f_c$ be a regular sequence in Q . Set $A = Q/(\mathbf{f})$. Let M be a finitely generated A -module with $\text{projdim}_Q M$ finite. Let I an ideal in A and let $N = \bigoplus_{n \geq 0} N_n$ be a finitely generated $\mathcal{R}(I)$ -module. Then*

$$\bigcup_{n \geq 0} \bigcup_{i \geq 0} \text{Ass Ext}_A^i(M, N_n) \text{ is a finite set.}$$

Furthermore there exists i_0, n_0 such that for all $i \geq i_0$ and $n \geq n_0$ we have

$$\begin{aligned} \text{Ass Ext}_A^{2i}(M, N_n) &= \text{Ass Ext}_A^{2i_0}(M, N_{n_0}) \\ \text{Ass Ext}_A^{2i+1}(M, N_n) &= \text{Ass Ext}_A^{2i_0+1}(M, N_{n_0}) \end{aligned}$$

The following example shows that two set of stable values of associate primes can occur

Example 5.2. Let $Q = k[[u, x]]$, $A = Q/(ux)$. Let $M = Q/(u)$, $I = A$ and $N = M[t]$ (so $N_n = M$ for all n).

For $i \geq 1$ one has

$$\text{Ext}_A^{2i-1}(M, M) = 0 \quad \text{Ext}_A^{2i}(M, M) = k.$$

5.3. We now state special case of a result due to E. West [14, 3.2;5.1].

Let $R = A[x_1, \dots, x_r; y_1, \dots, y_s]$ be a bi-graded A -algebra with $\deg x_i = (2, 0)$ and $\deg y_j = (0, 1)$. Let $M = \bigoplus_{i, n \geq 0} M_{(i, n)}$ be a finitely generated R -module. Then

- (1) $\bigcup_{i \geq 0} \bigcup_{n \geq 0} \text{Ass}_A M_{(i, n)}$ is a finite set.
- (2) $\exists i_0, n_0$ such that for all $i \geq i_0$ and $n \geq n_0$ we have

$$\begin{aligned} \text{Ass}_A M_{(2i, n)} &= \text{Ass}_A M_{(2i_0, n_0)} \\ \text{Ass}_A M_{(2i+1, n)} &= \text{Ass}_A M_{(2i_0+1, n_0)} \end{aligned}$$

First proof of Theorem 5.1. The result follows from our main Theorem 1 and 5.3. $\square$

For the convenience of the readers we give a self-contained second proof of Theorem 5.1. We need the following exercise problem from Matsumara's text (6.7, page 42) [9].

Fact 5.4. *Let $f: A \rightarrow B$ be a ring homomorphism of Noetherian rings. Let U be a finitely generated B -module. Then*

$$\text{Ass}_A U = \{\mathfrak{P} \cap A \mid \mathfrak{P} \in \text{Ass}_B U\}$$

In particular $\text{Ass}_A U$ is a finite set.

We will also need the following

Lemma 5.5. *Let $R = A[x_1, \dots, x_r; y_1, \dots, y_s]$ be a bi-graded A -algebra with $\deg x_i = (1, 0)$ and $\deg y_j = (0, 2)$. Let $D = \bigoplus_{i, j \geq 0} D_{(i, j)}$ be a finitely generated R -module. Set $l = r \bullet s$. Then there exists i_0, j_0 such that there exists inclusions*

$$D_{(i, j)} \hookrightarrow D_{(i+1, j+2)}^l \quad \text{for all } i \geq i_0 \text{ \& } j \geq j_0$$

Proof. Let $R_{++} = \bigoplus_{i \geq 1, j \geq 1} R_{(i,j)}$ be the irrelevant ideal of R . Set

$$E = D/H_{R_{++}}^0(D).$$

It is easy to check that $D_{(i,j)} = E_{(i,j)}$ for all $i, j \gg 0$. So if $E = 0$ then we have nothing to prove.

We consider the case when $E \neq 0$. Notice that

$$H_{R_{++}}^0(E) = 0. \quad \text{So } \text{grade}(R_{++}, E) > 0.$$

Now R_{++} is generated by the l elements $x_i y_j$; where $1 \leq i \leq r$ and $1 \leq j \leq s$. Notice also that $\deg x_i y_j = (1, 2)$ for all i, j . Consider the map

$$\begin{aligned} \phi: E &\mapsto (E(1, 2))^l \\ d &\mapsto x_i y_j d. \end{aligned}$$

Since $\text{grade}(R_{++}, E) > 0$ we get that ϕ is injective. Since $E_{(i,j)} = D_{(i,j)}$ for all $i, j \gg 0$; the result follows. $\square$

Second proof of Theorem 5.1. Let $\mathcal{S} = \mathcal{R}(I)[t_1, \dots, t_c]$. Then $D = E(N)$ is a finitely generated $\mathcal{S}$ -module.

1. By 5.4, it follows that $\text{Ass}_A E(N)$ is a finite set. Notice that

$$\text{Ass}_A \mathcal{E}(N) = \bigcup_{n \geq 0} \bigcup_{i \geq 0} \text{Ass } \text{Ext}_A^i(M, N_n).$$

2. By 5.5 it follows that we have inclusions

$$\text{Ext}_A^i(M, N_n) \hookrightarrow \text{Ext}_A^{i+2}(M, N_{n+1}) \quad \text{for all } i, n \gg 0.$$

It follows that

$$\text{Ass}_A \text{Ext}_A^i(M, N_n) \subseteq \text{Ass}_A \text{Ext}_A^{i+2}(M, N_{n+1}) \quad \text{for all } i, n \gg 0.$$

Using 1. we get the result. $\square$

To prove an analog of Theorem 5.1 for a local complete intersection we need the following result.

Lemma 5.6. *Let $(A, \mathfrak{m})$ be a Noetherian local ring. Let $\widehat{A}$ be the completion of A with respect to $\mathfrak{m}$. Let B be a finitely generated $\widehat{A}$ -algebra containing $\widehat{A}$. Let E be an A -module such that $E \otimes_A \widehat{A}$ is a finitely generated B -module. Let D be any finitely generated A -module. Then*

- (a) $\text{Ass}_{\widehat{A}} E \otimes_A \widehat{A}$ is a finite set.
- (b) $\text{Ass}_A E \otimes_A \widehat{A}$ is a finite set.
- (c) $\text{Ass}_A E = \text{Ass}_A E \otimes_A \widehat{A}$. In particular $\text{Ass}_A E$ is a finite set.
- (d) $\text{Ass}_A D = \text{Ass}_A D \otimes_A \widehat{A}$.

To prove this result we need Theorem 23.3 from [9]. Unfortunately there is a typographical error in the statement of Theorem 23.3 in [9]. So we state it here.

Theorem 5.7. *Let $\varphi: A \rightarrow B$ be a homomorphism of Noetherian rings, and let E be an A -module and G a B -module. Suppose that G is flat over A ; then we have the following:*

- (i) if $\mathfrak{p} \in \text{Spec } A$ and $G/\mathfrak{p}G \neq 0$ then

$${}^a\varphi(\text{Ass}_B(G/\mathfrak{p}G)) = \text{Ass}_A(G/\mathfrak{p}G) = \{\mathfrak{p}\}.$$

$$(ii) \text{ Ass}_B(E \otimes G) = \bigcup_{\mathfrak{p} \in \text{Ass}_A(E)} \text{Ass}_B(G/\mathfrak{p}G).$$

Remark 5.8. In [9] $\text{Ass}_A(E \otimes G)$ is typed instead of $\text{Ass}_B(E \otimes G)$.

Proof of Theorem 5.6. We consider the natural ring homomorphisms

$$\begin{aligned} \alpha: A &\hookrightarrow \widehat{A} \\ \beta: \widehat{A} &\hookrightarrow B \\ \gamma: A &\hookrightarrow B \end{aligned}$$

Clearly $\gamma = \beta \circ \alpha$.

- (a) We use the map β and fact 5.4 to get our result.
- (b) We use the map γ and fact 5.4 to get our result.
- (c) On the spectrum of the rings we have:

$${}^a\gamma = {}^a\alpha \circ {}^a\beta.$$

Therefore using fact 5.4 we get

$$\begin{aligned} \text{Ass}_A E \otimes \widehat{A} &= {}^a\alpha(\text{Ass}_{\widehat{A}} E \otimes \widehat{A}) \\ &= \{P \cap A \mid P \in \text{Ass}_{\widehat{A}} E \otimes \widehat{A}\}. \end{aligned}$$

We consider the flat extension $\alpha: A \rightarrow \widehat{A}$. By Theorem 5.7.(ii) (with $G = \widehat{A}$) we have

$$(*) \quad \text{Ass}_{\widehat{A}}(E \otimes_A \widehat{A}) = \bigcup_{\mathfrak{P} \in \text{Ass}_A E} \text{Ass}_{\widehat{A}} \widehat{A}/\mathfrak{P}\widehat{A}.$$

Now $\widehat{A}$ is faithfully flat. So if $\mathfrak{P} \in \text{Spec } A$ then $\widehat{A}/\mathfrak{P}\widehat{A} \neq 0$. By Theorem 5.7.(i) we get

$${}^a\alpha \left(\text{Ass}_{\widehat{A}} \frac{\widehat{A}}{\mathfrak{P}\widehat{A}} \right) = \mathfrak{P}$$

Taking ${}^a\alpha$ on (*) yields the desired result.

- (d) Set $B = \widehat{A}$. The result follows from (c). □

We now prove the following:

Theorem 5.9. *Let $(A, \mathfrak{m})$ be a local complete intersection. Let M be a finitely generated A -module. Let I an ideal in A and let $N = \bigoplus_{n \geq 0} N_n$ be a finitely generated $\mathcal{R}(I)$ -module. Then*

$$\bigcup_{n \geq 0} \bigcup_{i \geq 0} \text{Ass Ext}_A^i(M, N_n) \text{ is a finite set.}$$

Furthermore there exists i_0, n_0 such that for all $i \geq i_0$ and $n \geq n_0$ we have

$$\begin{aligned} \text{Ass Ext}_A^{2i}(M, N_n) &= \text{Ass Ext}_A^{2i_0}(M, N_{n_0}) \\ \text{Ass Ext}_A^{2i+1}(M, N_n) &= \text{Ass Ext}_A^{2i_0+1}(M, N_{n_0}) \end{aligned}$$

Proof. We consider the flat extension $\alpha: A \rightarrow \widehat{A}$. Say $\widehat{A} = Q/(f_1, \dots, f_c)$ where $(Q, \mathfrak{n})$ is a regular local ring and $f_1, \dots, f_c \in \mathfrak{n}^2$ is a regular sequence. 1. Consider $\mathcal{E}(N) = \bigoplus_{i \geq 0} \bigoplus_{n \geq 0} \text{Ext}_A^i(M, N_n)$ as an A -module. By Theorem 1; $E \otimes \widehat{A}$ is a

finitely generated $B = \mathcal{R}(\widehat{IA})[t_1, \dots, t_c]$ -algebra. By Lemma 5.6 we get that $\text{Ass}_A E$ is a finite set. Notice

$$\text{Ass}_A \mathcal{E}(N) = \bigcup_{n \geq 0} \bigcup_{i \geq 0} \text{Ass}_A \text{Ext}_A^i(M, N_n)$$

2. Set $\mathcal{E} = \mathcal{E}(N)$. By 5.5 we have injective maps

$$(*) \quad E_{(i,n)} \otimes \widehat{A} \hookrightarrow \left(E_{(i+1,n+2)} \otimes \widehat{A} \right)^l \quad \text{for all } i, n \gg 0.$$

Using Lemma 5.6 with $B = \widehat{A}$ and $E = E_{(i,n)}$ we get that for each $i \geq 0$ and $n \geq 0$

$$\text{Ass}_A E_{(i,n)} = \text{Ass}_A E_{(i,n)} \otimes_A \widehat{A}$$

The result follows from 1. and (*). $\square$

6. APPLICATION II SUPPORT VARIETIES

Let $(A, \mathfrak{m})$ be a local complete intersection of codimension c . Let M, N be two finitely generated A -modules. Define

$$\text{cx}_A(M, N) = \inf \left\{ b \in \mathbb{N} \mid \overline{\lim}_{n \rightarrow \infty} \frac{\mu(\text{Ext}_A^n(M, N))}{n^{b-1}} < \infty \right\}$$

In this section we prove the following theorem

Theorem 6.1. *Let $(A, \mathfrak{m})$ be a local complete intersection, M, N two finitely generated A -modules and let I be a proper ideal in A . Then*

$$\text{cx}_A(M, I^j N) \quad \text{is constant for all } j \gg 0.$$

6.2. Reduction to the case when A is complete and residue field of A is algebraically closed

6.3. Suppose A' is a flat local extension of A such that $\mathfrak{m}' = \mathfrak{m}A'$ is the maximal ideal of A' . If E is an A -module then set $E' = E \otimes_A A'$. Notice $I' \cong IA'$ and we consider it as an ideal in A' . By [1, 7.4.3] A' is also a complete intersection. It can be easily checked that

$$\text{cx}_{A'}(M', (I')^j N') = \text{cx}_A(M, I^j N) \quad \text{for all } n \geq 0.$$

We now do our reduction in two steps

By [3, App. Théorème 1, Corollaire], there exists a flat local extension $A \subseteq \widetilde{A}$ such that $\widetilde{\mathfrak{m}} = \mathfrak{m}\widetilde{A}$ is the maximal ideal of $\widetilde{A}$ and the residue field $\widetilde{k}$ of $\widetilde{A}$ is an algebraically closed extension of k . By 6.3 it follows that we may assume k to be algebraically closed. We now complete A . Note that $\widehat{A}$ is a flat extension of A which satisfies 6.3.

Thus we may assume that our local complete intersection A

- (1) is complete. So $A = Q/(f_1, \dots, f_c)$ where $(Q, \mathfrak{n})$ is regular local and $f_1, \dots, f_c \in \mathfrak{n}^2$ is a regular sequence.
- (2) The residue field of k is algebraically closed.

Of course there exists many Q and $f_1, \dots, f_c$ of the type as indicated above. We simply fix one such representation of A

6.4. Let U, V be two finitely generated A -modules.

Let $\mathcal{E}(U, V) = \bigoplus_{n \geq 0} \text{Ext}_A^n(U, V)$ be the total ext module of U and V . We consider it as a (finitely generated) module over the ring of cohomological operators $A[t_1, \dots, t_c]$. Since $\text{projdim}_Q U$ is finite $\mathcal{E}^*(U, V)$ is a finitely generated $A[t_1, \dots, t_c]$ -module.

6.5. Let $\mathcal{C}(U, V) = \mathcal{E}(U, V) \otimes_A k$. Clearly $\mathcal{C}(U, V)$ is a finitely generated $T = k[t_1, \dots, t_c]$ -module. (Here degree of t_i is 2 for each $i = 1, \dots, c$). Set

$$\mathfrak{a}(U, V) = \text{ann}_T \mathcal{C}(U, V).$$

Notice that $\mathfrak{a}(U, V)$ is a homogeneous ideal.

6.6. We now forget the grading of T and consider the affine space $\mathbb{A}^c(k)$. Let

$$\mathcal{V}(U, V) = \mathcal{V}(\mathfrak{a}(U, V)) \subseteq \mathbb{A}^c(k).$$

Since $\mathfrak{a}(U, V)$ is graded ideal we get that $\mathcal{V}(U, V)$ is a cone.

6.7. By a result due to Avramov and Buchweitz [2, 2.4] we get that

$$\dim \mathcal{V}(U, V) = \text{cx}_A(M, N)$$

Lemma 6.8. *If I is an ideal in A then there exists $j_0 \geq 0$ such that*

$$\mathcal{V}(U, I^j V) = \mathcal{V}(U, I^{j_0} V) \quad \text{for all } j \geq j_0.$$

We give a proof of Theorem 6.1 assuming Lemma 6.8.

Proof of Theorem 6.1. By 6.3 we may assume that A is also complete and has an algebraically closed residue field. The result now follows from 6.7 and 6.8. $\square$

6.9. Let $N = \bigoplus_{j \geq 0} I^j V$. Set $M = U$. By hypothesis $A = Q/(f_1, \dots, f_c)$ where $(Q, \mathfrak{n})$ is a regular local ring and $f_1, \dots, f_c \subseteq \mathfrak{n}^2$ is a regular sequence.

6.10. Set $\mathcal{C}(N) = \mathcal{E}(N) \otimes_A k$. By Theorem 1 $\mathcal{E}(N)$ is a finitely generated $\mathcal{S} = \mathcal{R}(I)[t_1, \dots, t_c]$ -module. It follows that $\mathcal{C}(N)$ is a finitely generated, bi-graded, $G = F(I)[t_1, \dots, t_c]$ -module. Recall that $F(I)$, the fiber-cone of I , is a finitely generated k -algebra. So we may as well consider $\mathcal{C}(N)$ as a bi-graded

$R = k[X_1, \dots, X_m, t_1, \dots, t_c]$ -module (of course here $X_1, \dots, X_m$ are variables). Furthermore $\deg X_l = (1, 0)$ for $l = 1, \dots, m$ and $\deg t_s = (0, 2)$ for $s = 1, \dots, c$. Set $T = k[t_1, \dots, t_c]$.

6.11. Advantages of coarsening the grading on $\mathcal{C}(N)$

By forgetting the degree on t 's we may consider $R = T[X_1, \dots, X_m]$. Notice that the correspondingly we obtain

$$\mathcal{C}(N) = \bigoplus_{n \geq 0} \mathcal{C}(U, I^n V)$$

We now give a

Proof of Lemma 6.8. We make the constructions as in 6.11. So $\mathcal{C}(N)$ is a finitely generated $R = T[X_1, \dots, X_m]$ -module. Notice that R is $\mathbb{N}$ -standard graded. So there exists j_0 such that

$$\text{ann}_T \mathcal{C}(N)_j = \text{ann}_T \mathcal{C}(N)_{j_0} \quad \text{for all } j \geq 0.$$

The results follows. $\square$

Question 6.12 (with hypothesis as in 6.1). *Is*

$$\mathrm{cx}_A(M, N/I^j N) \quad \text{constant for all } j \gg 0?$$

REFERENCES

- [1] L. L. Avramov, *Infinite free resolutions*, Six lectures on commutative algebra (Bellaterra, 1996), Progr. Math., vol. 166, Birkhäuser, Basel, 1998, pp. 345–392. MR MR1648669 (99j:13012)
- [2] Luchezar L. Avramov and Ragnar-Olaf Buchweitz, *Support varieties and cohomology over complete intersections*, Invent. Math. **142** (2000), no. 2, 285–318. MR MR1794064 (2001j:13017)
- [3] N. Bourbaki, *Algebré commutative. ix: Anneaux locaux réguliers complets*, Masson, Paris, 1983.
- [4] M. Brodmann, *Asymptotic stability of $\mathrm{Ass}(M/I^n M)$* , Proc. Amer. Math. Soc. **74** (1979), no. 1, 16–18. MR MR521865 (80c:13012)
- [5] Winfried Bruns and Jürgen Herzog, *Cohen-Macaulay rings*, Cambridge Studies in Advanced Mathematics, vol. 39, Cambridge University Press, Cambridge, 1993. MR MR1251956 (95h:13020)
- [6] David Eisenbud, *Homological algebra on a complete intersection, with an application to group representations*, Trans. Amer. Math. Soc. **260** (1980), no. 1, 35–64. MR MR570778 (82d:13013)
- [7] Tor H. Gulliksen, *A change of ring theorem with applications to Poincaré series and intersection multiplicity*, Math. Scand. **34** (1974), 167–183. MR MR0364232 (51 #487)
- [8] Daniel Katz and Eric West, *A linear function associated to asymptotic prime divisors*, Proc. Amer. Math. Soc. **132** (2004), no. 6, 1589–1597 (electronic). MR MR2051118 (2005d:13024)
- [9] Hideyuki Matsumura, *Commutative ring theory*, second ed., Cambridge Studies in Advanced Mathematics, vol. 8, Cambridge University Press, Cambridge, 1989, Translated from the Japanese by M. Reid. MR MR1011461 (90i:13001)
- [10] Leif Melkersson and Peter Schenzel, *Asymptotic prime ideals related to derived functors*, Proc. Amer. Math. Soc. **117** (1993), no. 4, 935–938. MR MR1124148 (93e:13027)
- [11] Tony J. Puthenpurakal, *The Hilbert function of a maximal Cohen-Macaulay module*, Math. Z. **251** (2005), no. 3, 551–573. MR MR2190344 (2006i:13022)
- [12] Anurag K. Singh, *p -torsion elements in local cohomology modules*, Math. Res. Lett. **7** (2000), no. 2-3, 165–176. MR MR1764314 (2001g:13039)
- [13] Wolmer V. Vasconcelos, *Cohomological degrees of graded modules*, Six lectures on commutative algebra (Bellaterra, 1996), Progr. Math., vol. 166, Birkhäuser, Basel, 1998, pp. 345–392. MR MR1648669 (99j:13012)
- [14] Eric West, *Primes associated to multigraded modules*, J. Algebra **271** (2004), no. 2, 427–453. MR MR2024817 (2004m:13004)

DEPARTMENT OF MATHEMATICS, INDIAN INSTITUTE OF TECHNOLOGY BOMBAY, POWAI, MUMBAI 400 076

E-mail address: tputhen@math.iitb.ac.in